

Technical Information

Phenolphthalein Phosphate Agar

Product Code: DM 1652

Application: - Phenolphthalein Phosphate Agar is recommended for identification of phosphatase positive *Staphylococcus aureus*.

Composition**

Ingredients	Gms / Litre
Peptic digest of animal tissue	5.000
Beef extract	3.000
Sodium chloride	5.000
Sodium phenolphthalein phosphate	0.012
Agar	15.000
Final pH (at 25°C)	7.4±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Staphylococcus are pathogens of man and other mammals. Depending on their ability to clot blood plasma (the coagulase reaction) they are divided into two groups. The coagulase-positive *staphylococci* constitute the most pathogenic species of *Staphylococcus aureus*. The presence of *staphylococci* in a lesion might first be suspected after examination of a direct gram stain. However, small numbers of bacteria in blood preclude microscopic examination and require culturing first ⁽¹⁾. Phosphatase has been implicated as a virulence factor for *S. aureus*. The organisms produce both an acid and alkaline phosphates, the latter being repressed in the presence of inorganic phosphate in the medium.

Phenolphthalein Phosphate Agar is used for the identification of phosphatase-positive colonies of *S. aureus*, which is a coagulase-positive pathogenic strain ⁽²⁾.

Peptic digest of animal tissue and beef extract supply the nitrogenous compounds, growth factors and trace ingredients essential for the growth of *Staphylococcus aureus*. Sodium phenolphthalein phosphate serves as a substrate for the phosphatase enzyme. Sodium chloride maintains osmotic equilibrium. Phosphatase production is determined by the liberation of phenolphthalein, which is indicated by the change in colour of the medium ⁽³⁾. When alkali is added to this medium, the liberated phenolphthalein gives a bright pink-red colouration ⁽⁴⁾. Alternatively phosphatase production can be determined by following technique.

Technique: Grow *staphylococci* overnight at 37°C on the medium. Invert the plate and pour few drops of ammonia solution into the lid, read a positive culture whose colonies turn bright pink within a few minutes. The colour soon fades.

Methodology

Suspend 28.01 grams of powder media in 1000 ml distilled water. Shake well & heat to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and dispense as desired.

Quality Control

Physical Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Yellow coloured clear to slightly opalescent gel forms in tubes as slants

Reaction

Reaction of 2.8% w/v aqueous solution at 25°C. pH : 7.4±0.2

pH Range 7.20-7.60



Dehydrated Culture Media
Bases / Media Supplements

Cultural Response/ characteristics

DM 1652: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum (CFU)	Growth	Recovery
Escherichia coli ATCC 25922	50-100	luxuriant	negative, no bright pink colour on addition of alkali
Staphylococcus aureus ATCC 25923	50-100	luxuriant	positive, bright pink colour on addition of alkali
Staphylococcus epidermidis ATCC 12228	50-100	luxuriant	positive, bright pink colour on addition of alkali

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Easmon C. S. F., Adlam C., 1983, Staphylococci and staphylococcal infections. Vol. 1 and 2, Academic Press, London,
2. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. 1, Williams and Wilkins, Baltimore.
3. Lewis B., 1961, J. Med. Lab. Technol., 18 : 112.
4. Barber M. and Kuper S. W. A., 1951, J. Pathol. Bacteriol., 63:65.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Donot use the products if it fails to meet specificatons for identity and performens parameters.

