



Dehydrated Culture Media
Bases / Media Supplements

Technical Information

Yeast Mannitol Agar w/ 1.5% Agar

Product Code: DM 1715

Application: Yeast Mannitol Agar w/ 1.5% Agar is used for cultivation, isolation and enumeration of soil microorganisms like *Rhizobium* species.

Composition**

Ingredients	Gms / Litre
Yeast extract	1.000
Mannitol	10.000
Dipotassium phosphate	0.500
Magnesium sulphate	0.200
Sodium chloride	0.100
Calcium carbonate	1.000
Agar	15.000
Final pH (at 25°C)	6.8±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Beijerinck was first to isolate and cultivate an aerobic gram negative rod-shaped microorganism from the nodules of legume. He named it *Bacillus radicolica*, which was subsequently placed under the genus *Rhizobium*. Bacteria belonging to the genus *Rhizobium* live freely in soil and in the root region of both leguminous and non-leguminous plants. However they show symbiosis only with leguminous plants by infecting their roots and forming nodules on them. *Rhizobium* present in the root nodules fixes atmospheric nitrogen i.e. gaseous nitrogen from air to organic nitrogen compounds, which is absorbed by plants. Thus role of *Rhizobium* is noteworthy for their major contributions to soil fertility. Yeast Mannitol Agar is used for the cultivation of symbiotic nitrogen fixing organism's viz. *Rhizobium* species⁽¹⁾.

Yeast extract serves as a good source of readily available amino acids, vitamin B complex and accessory growth factors for Rhizobia. It also poises oxidation - reduction potential of medium in the range favorable for Rhizobia and serves as hydrogen donor in respiratory process⁽²⁾.

Mannitol is the fermentable sugar alcohol source. Calcium and magnesium provide cations essential for the growth of Rhizobia.

Methodology

Suspend 27.8 grams of powder media in 1000 ml distilled water. Shake well & heat just to boiling. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and pour into sterile Petri plates.

Note: Due to presence of calcium carbonate, the prepared medium forms opalescent solution with white precipitate.



Quality Control

Physical Appearance

White to cream homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel.

Colour and Clarity of prepared medium

Whitish buff coloured opalescent gel forms in Petri plates.

Reaction

Reaction of 2.78% w/v aqueous solution at 25°C. pH : 6.8±0.2

pH range

6.60-7.00

Cultural Response

DM 1715: Cultural characteristics observed after an incubation at 25-30°C for upto 5 days.

Organism	Growth
<i>Rhizobium leguminosarum</i> ATCC 10004	luxuriant
<i>Rhizobium meliloti</i> ATCC 9930	luxuriant

Storage & Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Subba Rao N.S., 1977, Soil Microorganisms and Plant Growth, Oxford and IBG Publishing Company..
2. Allen. E.K. and Allen. O.N., 1950, Bacteriol. Rev., 14:273.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Donot use the products if it fails to meet specificatons for identity and performens parameters.