

Technical Information

Casman Broth Base

Product Code: DM 1766

Application: - Casman Broth Base with blood is used for isolation of fastidious microorganisms from clinical specimens under reduced oxygen tension.

Composition**

Ingredients	Gms / Litre
Proteose peptone	10.000
Tryptose	10.000
Beef extract	3.000
Dextrose	0.500
Corn starch	1.000
Sodium chloride	5.000
Nicotinamide	0.050
p-Amino benzoic acid (PABA)	0.050
Final pH (at 25°C)	7.2±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

For in vitro cultivation of fastidious microorganisms such as *Haemophilus* and *Neisseria* require the addition of X and V- growth factors Casman⁽¹⁻³⁾ described a blood-enriched medium for cultivation of *Haemophilus* and gonococci⁽¹⁾. The medium was developed to replace the previously described formulation that was time-consuming preparations using fresh and heated blood and meat infusion to supply the essential nutrients for growth of these fastidious organisms^(2, 3). Blood supplies both factor-X (hemin) and factor-V (Nicotinamide Adenine Dinucleotide), which is required for growth of *Haemophilus influenzae*. Sheep blood lacks factor-V due to NADase, an enzyme that destroys factor-V⁽⁴⁾. Horse and rabbit blood supplies both the factor X and factor V, and are relatively free of NADase activity, therefore it is preferred over sheep blood. Nicotinamide is added to medium to inhibit nucleotidase of erythrocytes that may destroy factor V.

Proteose peptone, tryptose and beef extract provide amino acids and other complex nitrogenous nutrients. Dextrose improves growth of pathogenic cocci. Corn starch prevents fatty acids from inhibiting the growth of *Neisseria gonorrhoeae*, without interfering with haemolytic reaction. Corn starch also neutralizes the inhibitory action of dextrose. Inoculate the medium as soon as the specimen arrives at the laboratory.

Methodology

Suspend 29.6 grams of powder media in 1000 ml distilled water. Shake well & heat to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 50°C and aseptically add 0.15% v/v sterile waterlysed blood (water:blood :: 3:1) of 5% sterile blood. Mix well and dispense as desired.

Quality Control

Physical Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Basal medium: Yellow coloured clear to slightly opalescent solution. After addition of blood: Cherry red coloured opalescent solution in tubes

Reaction

Reaction of 2.96% w/v aqueous solution at 25°C. pH : 7.2±0.2

pH range 7.00-7.40

Cultural Response/ characteristics

DM 1766: Cultural characteristics observed with added water-lysed blood, after an incubation at 35-37°C for 40-48 hours.



Dehydrated Culture Media
Bases / Media Supplements

Organism	Inoculum (CFU)	Growth
Haemophilus influenzae ATCC 35056	50-100	good-luxuriant
Neisseria meningitidis ATCC 13090	50-100	luxuriant
Streptococcus mitis ATCC 9895	50-100	luxuriant
Streptococcus pneumoniae ATCC 6303	50-100	luxuriant
Streptococcus pyogenes ATCC 19615	50-100	luxuriant

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Casman, 1947, Am. J. Clin. Pathol., 17:28 1.
2. Casman, 1942, J. Bact., 43:33.
3. Casman, 1947, J. Bact., 53:561.
4. Krunveide and Kuttner, 1938, J. Exp. Med., 67:429.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.

