



Dehydrated Culture Media  
Bases / Media Supplements

## Technical Information

### Maintenance (SCY) Medium

**Product Code: DM 1777**

**Application:** - Maintenance (SCY) Medium is used for the maintenance of iron bacteria.

#### Composition\*\*

Ingredients	Gms / Litre
Casein enzymic hydrolysate	0.910
Papaic digest of soyabean meal	0.030
Yeast extract	0.250
Sucrose	1.000
Sodium chloride	0.050
Dipotassium hydrogen phosphate	0.020
Thiamine	0.0004
Agar	10.000
Final pH (25°C)	7.3±0.2

\*\*Formula adjusted, standardized to suit performance parameters

#### Principle & Interpretation

Iron bacteria have the property of metabolizing reduced iron when present in their aqueous habitat and depositing it in the form of hydrated ferric oxide on or in their mucilaginous secretions. The large amount of brown slime so produced will impart a reddish tinge and an unpleasant odour to drinking water and make the supply unsuitable for domestic or industrial purposes.

Maintenance (SCY) Medium is prepared in accordance with APHA <sup>(1)</sup> and is used for the maintenance of iron bacteria. This media has proven successful for identifying various groups of filamentous organisms including iron bacteria <sup>(2)</sup>. Iron bacteria, especially those belonging to Sphaerotilus-Leptothrix group thrive in this media, which is too dilute to support proliferation of more rapidly growing organisms. Casein enzymic hydrolysate, yeast extract, papaic digest of soyabean meal and thiamine in the medium provide the necessary carbon, nitrogen, vitamins and minerals. Sucrose is the carbon source. Dipotassium phosphate provides buffering to the medium and sodium chloride provides the essential ions. Prepare agar slants and aseptically pipette 3 ml sterile tap water on the slant surfaces. Inoculate and incubate at room temperature until turbid growth develops in liquid layer. The cells remain viable for 3 months at refrigeration temperature.

#### Methodology

Suspend 12.26 grams of powder media in 1000 ml distilled water. Shake well & heat to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Aseptically add filter-sterilized solution of cyanocobalamin to a final concentration of 0.01 mg/litre. Mix well and dispense as desired.

#### Quality Control

##### Physical Appearance

White to cream homogeneous free flowing powder

##### Gelling

Firm, comparable with 1.0% Agar gel



**Colour and Clarity of prepared medium**

Cream coloured clear to slightly opalescent gel forms in Petri plates or in tubes as slants

**Reaction**

Reaction of 1.2% w/v aqueous solution at 25°C, pH: -7.3±0.2

**pH range** 7.10-7.50

**Cultural Response/ characteristics**

DM 1777: Cultural characteristics observed after an incubation at 25-30°C for 48-72 hours with added cyanocobalamin.

**Organism**

Leptothrix discophora ATCC 43182

Sphaerotilus natans ATCC 13338

Thiobacillus thioparus ATCC 8158

**Growth**

luxuriant

luxuriant

luxuriant

**Storage and Shelf Life**

**Dried Media:** Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

**Prepared Media:** 2-8° in sealable plastic bags for 2-5 days.

**Further Reading**

1. Eaton A. D., Clesceri L. S., Rice E. W. and Greenberg A W.(Eds.), 2005, Standard Methods for the Examination of Water and Wastewater, 21st Ed., APHA, Washington, D.C.
2. Van Veen W. L., 1973, Antonie Van Leeuwenhoek (Holland), 39:189.

**Disclaimer :**

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.