

## Technical Information

### TN Agar

**Product Code: DM 1950**

**Application:** - TN Agar is used for isolation and cultivation of *Vibrios* from food samples.

### Composition\*\*

Ingredients	Gms / Litre
Casein enzymic hydrolysate	10.000
Sodium chloride	10.000
Agar	15.000
Final pH ( at 25°C)	7.2±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Principle & Interpretation

Members of the genus *Vibrio* are defined as gram-negative, asporogenous rods that are straight or have a single rigid curve. Three species of *Vibrio*, namely *Vibrio cholerae*, *Vibrio parahaemolyticus* and *Vibrio mimicus* are well-documented as human pathogens.

*V. cholerae*, the type species of the genus *Vibrio* is the causative agent of cholera outbreaks and epidemics.

*V. cholerae* can be differentiated from other *Vibrio* species except *V. mimicus*, because of its obligate requirement for sodium ions (Na<sup>+</sup>). TN Agar is formulated according to APHA <sup>(1)</sup> for cultivation of *Vibrio* species from foods.

Casein enzymic hydrolysate provides nitrogenous, carbonaceous compounds, sulphur, vitamin B complex and other essential growth nutrients. Sodium chloride improves the selectivity of the medium.

Weigh 25 grams of food sample such as seafood or vegetable and blend or cut into small pieces into two jars. To one jar add 225 ml Alkaline Peptone Water (DM1618) and to another jar add 225 ml Gelatin Phosphate Salt Broth. Incubate at 35 ± 2°C for 6 to 8 hours. Transfer a loopful from surface growth of each broth culture to two plated media, i.e. TCBS Agar (DM1189) and Gelatin Phosphate Salt Agar (DM1921), and incubate at 35 ± 2°C for 18-24 hours. Subculture 3-4 colonies from each plating medium to TN Agar. Growth from TN Agar is further confirmed by inoculating Kligler Iron Agar slants (DM1078) and stabbing the butt. After incubation, *V. cholerae* cultures will have an alkaline (red) slant and an acid (yellow) butt, no gas, and no blackening (H<sub>2</sub>S production) in the butt. Presumptive test for suspected strains of *V. cholerae* from TN Agar is carried out by using string test <sup>(2)</sup>.

### Methodology

Suspend 35 grams of powder media in 1000 ml distilled water. Shake well & heat to dissolve the medium completely. Dispense in tubes. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Allow the tubed medium to solidify in an inclined position (long slants).

### Quality Control

#### Physical Appearance

Cream to yellow homogeneous free flowing powder

#### Gelling

Firm, comparable with 1.5% Agar gel.

#### Colour and Clarity of prepared medium

Light yellow coloured clear to slightly opalescent gel forms in tubes as long slants.

#### Reaction

Reaction of 3.5% w/v aqueous solution at 25°C. pH : 7.2±0.2

#### pH range

7.00-7.40

#### Cultural Response/Characteristics

**DM 1950:** Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum(CFU)	Growth
<i>Vibrio cholerae</i> ATCC 15748	50-100	good-luxuriant
<i>Vibrio parahaemolyticus</i> ATCC 17802	50-100	good-luxuriant



Dehydrated Culture Media  
Bases / Media Supplements

## Storage and Shelf Life

**Dried Media:** Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

**Prepared Media:** 2-8<sup>0</sup> in sealable plastic bags for 2-5 days.

## Further Reading

1. Downes F. P. and Ito K., (Eds.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 4th Ed., APHA, Washington, D.C.
2. Smith H. L. Jr., 1970, Bull. World Health Organization, 42:817.

## Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.

