



Dehydrated Culture Media
Bases / Media Supplements

Technical Information

Lactic Bacteria Differential Broth

Product Code: DM 2086

Application: - Lactic Bacteria Differential Broth is used for differentiation of homofermentative and heterofermentative lactic acid bacteria.

Composition**

Ingredients	Gms / Litre
Casein enzymic hydrolysate	10.000
Papaic digest of soyabean meal	1.500
Casein acid hydrolysate	3.000
Yeast extract	1.000
Fructose	2.500
Monopotassium phosphate	2.500
Bromocresol green	0.055
Final pH (at 25°C)	7.0±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Lactic Bacteria Differential Broth is devised as per McDonald et al (1) for differentiation of homofermentative Lactobacilli and heterofermentative Streptococci. Lactobacilli and Streptococci are used as starter cultures in food and dairy industry. Streptococci grow first and produce metabolites, lowering redox potential which enables Lactobacilli to grow. Lactobacilli synthesize products which stimulate growth of Streptococci. Medium constituents like casein acid hydrolysates, papaic digest of soyabean meal and yeast extract supply all the necessary nutrients for the growth of lactic bacteria. Fructose is the fermentable carbohydrate in the medium. Bromo cresol green is the pH indicator. Heterofermentative lactic acid bacteria produce CO₂, lactic acid, acetic acid, ethanol and mannitol. Homofermentative lactic acid bacteria produce only lactic acid. Homofermentative lactic acid bacteria produce lactic acid from fructose and is indicated by yellow colour formation. Heterofermentative lactic acid bacteria induce lesser acidification and thus vary in the colour formation by the indicator in the medium

Methodology

Suspend 20.5 grams of powder media in 1000 ml distilled water. Add 1 gram of polysorbate 80. Shake well & heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and dispense as desired.

Quality Control

Physical Appearance

Light yellow to bluish grey homogeneous free flowing powder

Colour and Clarity of prepared medium

Blue coloured clear solution in tubes

Reaction

Reaction of 2.05% w/v aqueous solution at 25°C. pH : 7.0±0.2

pH Range 6.80-7.20





Dehydrated Culture Media
Bases / Media Supplements

Cultural Response/Characteristics

DM 2086: Cultural characteristics observed after an incubation at 35-37°C for 18-48 hours

Cultural Response	Inoculum (CFU)	Growth	Colours of medium
<i>Lactobacillus casei</i> ATCC 9595	50-100	Luxuriant	Green
<i>Lactobacillus plantarum</i> ATCC 8014	50-100	Luxuriant	Green
<i>Streptococcus cremoris</i> ATCC 19257	50-100	Luxuriant(incubated at 30 ^o C)	Blue
<i>Streptococcus thermophilus</i> ATCC 14485	50-100	Luxuriant(incubated at 45 ^o C)	Bluish-green

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8^o in sealable plastic bags for 2-5 days.

Further Reading

1. McDonald L.C., McFeters R.F., Daeschel M.A. and Fleming H.P., 1987, Appl. Environ. Microbiol., 53:1382.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.

