

Technical Information

Luria Bertani Agar Miller

Product Code : DM2151

Application: - Luria Bertani Agar Miller is used for the cultivation and maintenance of recombinant strains of *Escherichia coli* for genetic and molecular studies and may be used for routine cultivation of not particularly fastidious microorganisms.

Composition**

Ingredients	Gms / Litre
Casein enzymic hydrolysate	10.000
Yeast extract	5.000
Sodium chloride	10.000
Agar	15.000
Final pH (at 25°C)	7.5±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Luria Bertani Agar as described by Lennox ⁽¹⁾ is used for cultivation and maintenance of recombinant strains of *Escherichia coli*. Luria Bertani Agar, Miller ⁽²⁾ is slightly different as it contains double amount of sodium chloride. The media is nutritionally rich for the growth of pure cultures of recombinant strains. Strains derived from *Escherichia coli* K12 are deficient in Vitamin B synthesis are further modified by specific mutation to create auxotrophic strains those are unable to grow on nutritionally deficient media.

Casein enzymic hydrolysate provides peptides and peptones while Vitamin B complex is provided by yeast extract. Sodium chloride provides sodium ions for membrane transport and also maintains the osmotic equilibrium of the medium.

Methodology

Suspend 40 grams of powder media in 1000 ml distilled water. Shake well heat to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Dispense as desired. Mix well and and pour into sterile Petri plates.

Quality Control

Physical Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Yellow to amber coloured, clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 4.0% w/v aqueous solution at 25°C. pH : 7.5±0.2

pH range

7.30-7.70

Cultural Response/Characteristics

DM2151 Cultural characteristics observed after an incubation at 35-37°C for 18 - 24 hours.



Dehydrated Culture Media
Bases / Media Supplements

Organism	Inoculum (CFU)	Growth	Recovery
<i>Escherichia coli</i> ATCC 23724	50-100	luxuriant	>=70%
<i>Escherichia coli</i> ATCC 25922	50-10	Luxuriant	>=70%
<i>Escherichia coli</i> DH5 alpha MTCC 1652	50-100	Luxuriant	>=70%

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8⁰ in sealable plastic bags for 2-5 days.

Further Reading

1. Lennox E.S., 1955, Transduction of Linked Genetic Characters of the host by bacteriophage P1, *Virology*, 1:190.
2. Atlas R.M., 1993, *Handbook of Microbiological Media*, Ed. by Parks L., CRC Press, Inc.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performance parameters.

