

## Technical Information

### Nutrient Agar for Oxidase

#### Product Code: DM 2274

**Application:** - Nutrient Agar is used for confirmation of presence of oxidase in microorganisms in water.

#### Composition\*\*

Ingredients	Gms / Litre
Peptic digest of animal tissue	1.000
Meat extract	1.000
Sodium chloride	5.000
Agar	15.000
Final pH ( at 25°C)	7.3±0.2

\*\*Formula adjusted, standardized to suit performance parameters

#### Principle & Interpretation

Nutrient Agar for oxidase is recommended by APHA <sup>(1)</sup> & ISO committee <sup>(2)</sup> for differentiation of the coliform bacteria on the basis of presence of enzyme cytochrome oxidase. Cytochrome oxidase is a iron-containing porphyrin enzyme that participates in the electron transfer mechanisms and in the nitrate metabolic pathways of some bacteria. Although the test can be performed by flooding the agar surface of an inoculated plate with the reagent after incubation or with the help of oxidase reagent impregnated filter paper.

Peptic digest of animal tissue and meat extract provide nitrogenous compounds, carbon, sulphur and trace ingredients. Sodium chloride maintains osmotic equilibrium.

Nutrient Agar plates are streak inoculated to obtain isolated colonies. The isolated colony is used for oxidase testing on an impregnated filter paper. A dark purple colour that develops within 10 seconds is a positive oxidase test. Coliform bacteria are oxidase negative.

#### Medhlogy

Suspend 22 grams of powder media in 1000 ml distilled water. Shake well & heat to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and pour into sterile Petri plates.

#### Quality Control

##### Physical Appearance

Cream to yellow homogeneous free flowing powder

##### Gelling

Firm, comparable with 1.5% Agar gel

##### Colour and Clarity of prepared medium

Yellow coloured clear to slightly opalescent gel forms in Petri plates

##### Reaction

Reaction of 2.2% w/v aqueous solution at 25°C. pH : 7.3±0.2

pH Range 7.10-7.50

##### Cultural Response/ characteristics

DM 2274: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.



Dehydrated Culture Media  
Bases / Media Supplements

Organism	Growth	Oxidase
Aeromonas hydrophila ATCC 7966	luxuriant	positive reaction, deep purple blue colour develops within 10 seconds
Escherichia coli ATCC 25922	luxuriant	negative reaction
Enterobacter aerogenes ATCC 13048	luxuriant	negative reaction
Pseudomonas aeruginosa ATCC 27853	luxuriant	positive reaction, deep purple blue colour develops within 10 seconds
Vibrio cholerae ATCC 15748	luxuriant	positive reaction, deep purple blue colour develops within 10 seconds

## Storage and Shelf Life

**Dried Media:** Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

**Prepared Media:** 2-8° in sealable plastic bags for 2-5 days.

## Further Reading

1. Greenberg A. E., Clesceri L. S. and Eaton A. D., (Eds.), 1992, Standard Methods for the Examination of Water and Wastewater, 18th Ed., APHA, Maryland.
2. International Organization for Standardization (ISO), 1990, Draft, ISO/DIS 9308-1.

## Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.

