

Technical Information

Mannitol Selenite Broth (Twin Pack) (Selenite Mannitol Broth)

Product Code: DM 2534

Application: - Mannitol Selenite Broth is used for selective enrichment of Salmonellae from clinical materials.

Composition**

Ingredients	Gms / Litre
Part A	-
Peptic digest of animal tissue	5.000
Mannitol	4.000
Sodium phosphate	10.000
Part B	-
Sodium hydrogen selenite(Sodium biselenite)	4.000
Final pH (25°C)	7.1±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Selenite-containing media for the enrichment of Salmonella was first reported by Guth ⁽¹⁾. This medium was further modified by Leifson ⁽²⁾ for the enrichment and isolation of Salmonella from clinical specimens. Mannitol Selenite Broth described by Hobbs and Allison ⁽³⁾ is a selective enrichment medium, which is more or less similar to Leifson ⁽²⁾ enrichment medium, being used for the isolation of Salmonella Typhi and Salmonella Paratyphi B from clinical specimens. Mannitol Selenite Broth can also be used for the selective enrichment of Salmonella from water and foodstuffs.

Peptic digest of animal tissue provides amino acids and other nitrogenous substances to Salmonellae. Mannitol serves as fermentable carbohydrate, a sugar alcohol which also helps in maintaining a uniform pH alongwith sodium phosphate. Sodium phosphate also lessens the toxicity of selenite.

Do not incubate longer than 24 hours as the inhibitory effect of selenite is reduced after 6 - 12 hours incubation ⁽⁴⁾. Subculture broth from the upper third of the broth column to greater or lesser inhibitory selective medium such as MacConkey Agar (DM1081)

Methodology

Suspend 4.0 grams of Part B in 1000 ml distilled water. Add 19.0 grams of Part A. Warm to dissolve the medium completely. Distribute in sterile test tubes. Sterilize in a boiling water bath or free flowing steam for 10 minutes. DO NOT AUTOCLAVE. Excessive heating is detrimental. Discard the prepared medium if large amount of selenite is reduced (indicated by red precipitate at the bottom of the tube).

Caution: Sodium hydrogen selenite (Sodium biselenite) is very toxic, corrosive agent and causes teratogenicity. So it should be handled with great care. If there is contact, wash immediately with lot of water.

Quality Control

Physical Appearance

Part A : Cream to yellow homogeneous free flowing powder Part B : White to cream homogeneous free flowing powder

Colour and Clarity of prepared medium

Light yellow coloured clear to slightly opalescent solution of complete medium

Reaction

Reaction of 1.9% w/v of Part A + 0.4% w/v of Part B at 25°C. pH : 7.1±0.2

pH range 6.90-7.30

Cultural Response/ characteristics

DM 2534: Cultural characteristics observed when subcultured on MacConkey Agar (DM1081), after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum (CFU)	Recovery(increase in numbers)	Colour of colony
Escherichia coli ATCC 25922	50-100	little-none	pink with bile precipitate
Salmonella Enteritidis ATCC 13076	50-100	luxuriant	colourless
Salmonella Paratyphi B ATCC 8759	50-100	luxuriant	colourless
Salmonella Typhi ATCC 6539	50-100	luxuriant	colourless

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Guth F., 1916, Zentralbl. Bakteriol. Parasitenk. Indektionskr. Hyg. Abt. 77:487
2. Leifson E., 1936, Am. J. Hyg., 24(2):423.
3. Hobbs B. C. and Allison V. D., 1945, Mon. Bull. Min. Hlth. Publ. Hlth. Lab. Serv., 4:12.
4. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. 1, Williams and Wilkins, Baltimore

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.