



Dehydrated Culture Media  
Bases / Media Supplements

## Technical Information

### SBG Enrichment Broth (Twin Pack)

**Product Code: DM 2535**

**Application:** - SBG Enrichment Broth is used for selective enrichment of *Salmonella* species from clinical specimens.

### Composition\*\*

Ingredients	Gms / Litre
Part A	5.000
Peptic digest of animal tissue	5.000
Yeast extract	5.000
Mannitol	1.000
Sodium taurocholate	2.650
Dipotassium phosphate	1.020
Monopotassium phosphate	0.005
Brilliant green	-
Part B	4.000
Sodium hydrogen selenite	7.2±0.2

Final pH ( at 25°C)

\*\*Formula adjusted, standardized to suit performance parameters

### Principle & Interpretation

*Salmonella* are gram-negative, facultatively anaerobic, non-sporulating, motile rods in the family Enterobacteriaceae. They are widely distributed in animals affecting mainly the stomach and the intestines and difficult to differentiate biochemically from *Escherichia coli*. Leifson's Selenite Medium <sup>(1)</sup> and Kauffmann's Modified Tetrathionate Medium have been widely used as enrichment medium for the isolation of *Salmonella*. Selenite Medium used for enrichment of *Salmonella* inhibits *E. coli* but allows growth of *Proteus* and *Enterobacter*. To overcome this difficulty, Stokes and Osborne developed a more selective medium by adding brilliant green and sodium taurocholate to the Selenite Medium and found that it was superior to the Selenite Medium for isolating *Salmonella* in patients with gastroenteritis and similar diseases.

SBG (Selenite Brilliant Green) Enrichment Broth is prepared as per the formulation described by Stokes and Osborne <sup>(2)</sup> for selective enrichment of *Salmonella* from clinical specimens and egg products. Brilliant green and sodium selenite are neutralized by the egg constituents' rendering the medium non-selective therefore sulfapyridine is added to the medium for isolation of *Salmonella* from eggs <sup>(3)</sup>.

1 gram or 1 ml of test material is inoculated in 10 ml of the medium and incubated at 35-37°C for 18-24 hours. Following incubation, a loopful of the enriched culture is streaked on SS Agar (DM1108), MacConkey Agar (DM1081) or other plates for the isolation of *Salmonella*.

Peptic digest of animal tissue and yeast extract provide nitrogenous compounds, carbon, sulphur, vitamin B complex and trace elements necessary for the growth of organisms. Mannitol is the fermentable carbohydrate. Mannitol is utilized by *Salmonella* as an energy source, but it cannot be utilized by *Proteus*. Phosphates buffer the medium well. Brilliant green, sodium hydrogen selenite, sodium taurocholate inhibit the growth of gram-positive organisms and enteric organisms except *Salmonella* species. Whole egg and egg yolk reduces the selective properties of Selenite-Brilliant Green Enrichment. Addition of sulfapyridine restores the selective properties <sup>(3)</sup>. This medium cannot be used for the isolation of *Salmonella Typhi*, *Salmonella Paratyphi A*, and *Salmonella Pullorum*.

### Methodology

Suspend 4 grams of Part B of in 1000 ml distilled water. Add 19.67 grams of Part A. Mix well. Shake well & heat to boiling for 5 to 10 minutes. DO NOT AUTOCLAVE OR OVERHEAT. Dispense in sterile tubes. Add 0.5 g/l sodium sulfapyridine if desired. Caution: Sodium hydrogen selenite (Sodium biselenite) is very toxic, corrosive agent and causes teratogenicity. So it should be handled with great care. If there is contact with skin wash immediately with lot of water.



## Quality Control

### Physical Appearance

Part A : Cream to greenish yellow homogeneous free flowing powder Part B : White to cream homogeneous free flowing powder

### Colour and Clarity of prepared medium

Light green coloured clear to slightly opalescent solution

### Reaction

Reaction of 1.97% w/v of Part A + 0.4% w/v of Part B at 25°C. pH : 7.2±0.2

**pH Range:-** 7.00-7.40

### Cultural Response/Characteristics

DM 2535: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours when subcultured on MacConkey Agar (DM1081)

Organism	Inoculum (CFU)	Growth	Recovery (on M081)	Colour of colony (on M081)
<i>Salmonella Choleraesuis</i> ATCC 12011	50-100	luxuriant	>=50%	Colourless
<i>Salmonella Typhi</i> ATCC 6539	50-100	luxuriant	>=50%	Colourless
<i>Salmonella Typhimurium</i> ATCC 14028	50-100	luxuriant	>=50%	Colourless
<i>Enterobacter aerogenes</i> ATCC 13048	50-100	None-poor	>=50%	Pink to colourless
<i>Escherichia coli</i> ATCC 25922	50-100	None-poor	<=10%	Pink with bile precipitation

## Storage and Shelf Life

**Dried Media:** Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

**Prepared Media:** 2-8° in sealable plastic bags for 2-5 days.

## Further Reading

1. Leifson, 1955, Appl. Microbiol. 3:295
2. Stokes and Osborne, 1955, Appl. Microbiol., 3:217.
3. Osborne and Stokes, 1955, Appl. Microbiol., 3:295.

## Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.